



INFLUENCE OF NOVA AMI-CA FOLIAR FERTILIZATION ON YIELD, YIELD COMPONENTS AND QUALITY CHARACTERISTICS OF PEANUT (*Arachis hypogaea* L.)

Venkata narayana, P., Sambasiva rao, Y., Kiran kumar, A., Sreekanth, Y., Basanth, N.

Nova Innovation Centre, Nova Agri Tech Ltd., Plot No. 57, Hanuman Nagar, Chinnatokatta, New Bowenpally, Secunderabad, Telangana – 500 011

*Corresponding author email: cro@novaagritech.com

ABSTRACT

In order to investigate the effect of Nova ami-ca on yield, yield components of peanut (*Arachis hypogaea* L.) in 2018 an experiment was conducted in Randomized Block Design (RBD) with treatments T₁: Control (No application of any fertilizer), T₂: Recommended dose of fertilizers (RDF) with 2 splits at 45 and 75 DAS, T₃: T₂+ 5g/l Calcium nitrate at 45, 75 DAS, T₄: T₂ + 2.5ml/l of Nova ami-Ca at 45 and 75 DAS, T₅: 5g/l Calcium nitrate at 45, 75DAS, T₆: 2.5ml/l of Nova ami-Ca at 45 and 75 DAS. Results showed that application of T₄ had significant effect on plant height, number of branches per plant, number of filled pods, no of unfilled pods, 100 kernel weight, oil content and pod yield but application of T₃ resulted highest protein content. According to the results of the present study calcium is an important factor for increasing of yield and yield components of peanut.

KEY WORDS: Peanut, Calcium, Nova ami-ca, Yield; Yield components.

INTRODUCTION

Ground nut is an important oil, protein, food and feed legume crop grown in over 100 countries. It covered 26.7 million hectares area worldwide with a total production of 37.1 million MT. Though, the groundnut cultivation has been extended in almost all soils throughout the world, its nutrient requirement is high and like other crops it also requires all the macro- and micro-nutrients for its growth and development (Dwivedi 1988, Singh *et al.*, 1990, Adams and Hartzog 1991, Adams *et al.*, 1993 and Singh 1999). Calcium is an essential plant nutrient that plays a significant role in peanut seed development. According to the Department of Agriculture (DOA) 2006 the application of Calcium is important for proper kernel development in groundnut. It is a constituent of cell walls and involved in production of new growing points and root tips. It provides elasticity and expansion of cell walls, which keeps growing points away from becoming rigid and brittle. It acts as a base for neutralizing organic acids generated during the growing process and aids in carbohydrate translocation and nitrogen absorption. Indeed, calcium might be considered as bricks in plant assembly. Application of amino chelate fertilizers by farmers has increased during recent years (Souri and Yarahmadi, 2016). Amino chelate fertilizers are the latest novelties regarding plant nutrition in agriculture (Souri, 2015). They are among the new and modern formulae of fertilizers which are synthesized based on various amino acids. Application of amino chelates instead of simple routine fertilizers generally results in higher nutrient uptake efficiency (Souri and Yarahmadi, 2016; Ghasemi., 2014; Garcia *et al.*, 2011). In many cases, only the use of chelated forms of micronutrients could meet the plant's requirements for nutrients to provide healthy growth and

high yield achievements (Ghasemi *et al.*, 2013 and Souri, 2015).

MATERIALS & METHODS

This study was conducted in the *kharif*, 2018 at the Nova Agritech Ltd. Experimental farm, Kamareddy, Telangana, India. To investigate the performance of Nova amino chelated-calcium on yield components and quality parameters of Ground nut. The experiment was laid out in Randomized block design (RBD) with three replications. The groundnut cultivar kadiri-9 with duration 110-120 days was used for this study. The field was prepared by ploughing and leveling, divided into small plots of 25m² (5m × 5m) by raising bunds and the seeds of Ground nut were sown at 30×10 cm spacing in the furrows and covered with soil. The crop was grown under recommended package of practices and proper care was taken to protect it from weeds, insects, pests and diseases during entire cropping season. Data was recorded on established plants at 60 days after planting and at harvest stage. Plant height, no of branches per plant, no. of filled pods, no. of unfilled pods, 100 kernel weight, oil content by according to Tiwari *et al.* (1974), protein content according to by Lowry *et al.*, (1951) and pod yields were investigated.

Treatments

- T₁: Control (No application of any fertilizer).
- T₂: Recommended dose of fertilizers (RDF) with 2 splits at 45 and 75 DAS.
- T₃: T₂+ 5g/l Calcium nitrate at 45, 75DAS.
- T₄: T₂ + 2.5ml/l of Nova ami-Ca at 45 and 75 DAS.
- T₅: 5g/l Calcium nitrate at 45, 75DAS.
- T₆: 2.5ml/l of Nova ami-Ca at 45 and 75 DAS.

RESULTS & DISCUSSIONS

Plant height

Data on plant height of ground nut at 60 DAS and at harvest presented in table 1. The results of this study shows that significantly maximum plant height with 28.9 and 35.76 cm at 60 DAS and at harvest respectively recorded in response to T₄ (T₂ + 2.5ml/l of Nova ami-Ca at 45 and 75 DAS), while minimum plant height 22.4 and

28.43 cm at 60DAS and at harvest respectively recorded in T₁(Control). Calcium is a important plant nutrient that affects cell wall and membrane formation and plays a key role in plant growth, biomass production (Madani *et al.*, 2015). Calcium application increased plant height by activating enzymes for cell mitosis, division and elongation and height (Jones, 1999).

TABLE 1: Effect of calcium source Nova ami-ca on plant height and No. of branches

Treatments	Plant height (cm)		No. of branches/plant	
	60 DAS	At harvest	60 DAS	At harvest
T ₁	22.4	28.43	5.51	5.81
T ₂	24.8	31.90	6.05	6.57
T ₃	26.5	32.46	6.09	6.95
T ₄	28.9	35.76	6.35	7.41
T ₅	23.9	28.94	5.57	6.45
T ₆	24.4	30.10	5.78	6.54
SEm	0.8795	1.4572	0.1393	0.1157
CD 5%	2.7714	4.5916	0.4390	0.3647
CV%	6.0561	8.0702	4.0949	3.0261

Number of branches per plant⁻¹

The obtained results shows that significant differences were observed with more number of branches in T₄ at 60 DAS and at harvest and lower number of branches noticed T₁ at 60DAS and at harvest. The increase in number of

branches per plant might be due to the role of Ca in cell division, mitosis and carbohydrate metabolism (Davis *et al.*, 2003) application of calcium increased number of branches (Kamara *et al.*, 2011)

TABLE 2: Effect of Nova ami-ca on yield and yield components of peanut

Treatments	No. of filled pods	No. of un filled pods	100 kernel weight	Oil content %	Protein content %	Pod yield/ha
T ₁	2.6	1.44	23.83	44.56	14.76	1114
T ₂	13.88	1.52	30.89	48.18	25.50	1691
T ₃	16.50	1.61	32.56	47.37	29.30	1934
T ₄	19.18	1.03	34.44	49.45	29.20	2024
T ₅	11.44	1.35	28.65	45.01	25.26	1665
T ₆	13.55	1.11	29.22	46.86	27.37	1695
SEm	0.5611	0.0743	0.5114	0.5745	0.4072	11.41
CD 5%	1.7679	0.2341	1.6114	1.8102	1.2830	35.96
CV%	7.5619	9.5615	2.9588	2.1212	2.8073	1.1714




FIGURE 1: Yield Components in Peanut

Number of filled pods plant⁻¹

Number of filled pods per plant (table 2) differed significantly among all treatments. Maximum number of filled pods noticed in T₄ followed by T₃ and minimum number of filled pods observed in T₁ calcium application increased the number of two seeded pods and increased pod set Son *et al.* (1974). Calcium application had a positive effect on the number of filled pods per plant (Kamara *et al.*, 2011).

Number of unfilled pods plant⁻¹

Significantly higher number of unfilled pods observed in T₁ in comparison to other treatments and lower number of unfilled pods noticed in T₄. Application of Ca significantly decreased the number of unfilled pods in ground nut (Kogram *et al.*, 1999)

100 kernel weight

Application of Recommended dose of fertilizers (RDF) with 2 splits at 45 and 75 DAS + 2.5ml/l of Nova ami-Ca at 45 and 75 DAS was showed significant differences among all treatments with highest weight observed in T₄ and lowest was recorded in T₁. The weight of 100 kernel had a significant effect to apply calcium (Gashti, 2012). Application of calcium fertilizer had a positive effect on the 100 seed weight (Kamara *et al.*, 2011)

Oil content (%)

Oil content highest was recorded in T₄ which differed significantly with all the treatments except T₂ which shows parity with T₄. Application of calcium results significant effect on oil content of ground nut (Gashti, 2012). Results showed that increased calcium fertilization increased the vigour, germination and oil content of the seeds (Kamara *et al.*, 2017)

Protein content (%)

This study confirms that lowest protein content recorded in T₁ which differed significantly with all other treatments. Highest protein content was noticed in T₃ followed by T₄. Badiger *et al.* (1982) concluded that crude protein per cent and oil yield were improved by the application of calcium.

Pod yield (kg ha⁻¹)

This study reveals that T₄ significantly differed with all other treatments and lowest pod yields observed in T₁. Calcium fertilization is important for seed yields of ground nut (Kamara *et al.*, 2011).

REFERENCES

Adams, J.F. and Hartzog, D.L. (1991) Seed quality of runner peanuts as affected by gypsum and soil calcium. *J. Plant Nutrition*. 14: 841-851.

Adams, J.F., Hartzog, D.L and Nelson, D.B. (1993) Supplemental calcium application on yield, grade and seed quality of runner peanut. *Agron. J.* 85: 86-93.

Badiger, M.K., Reddy, N.P.S., Michael Rand Shivraj, B. (1982) Influence of fertilizer potassium, sulphur and calcium on the yield and quality attributes of groundnut. *Journal of Indian Society of Soil Science* 30: 166-169.

Davis, J.M., Sanders, D.C., Nelson, P.V., Lengnick, L. and Sperry, W.J. (2003) Boron improves growth, yield, quality, and nutrient content of tomato. *Journal of the American Society for Horticultural Science*, 128(3):441-446.

Department of Agriculture (2006) Annual report. Socio Economic & Planning Centre, Department of Agriculture.

Dwivedi, R.S. (1988) Mineral Nutrition of Groundnut.

Garcia, A.L., Madrid, R., Gimeno, V., Rodriguez-Ortega, W.M., Nicolas, N. and Garcia-Sanchez, F. (2011) The effects of amino acids fertilization incorporated to the nutrient solution on mineral composition and growth in tomato seedlings. *Spanish Journal of Agricultural Research*. 9 (3): 852-861.

Gashti, A.H., Vishekaei, M.N.S and Hosseinzadeh, M.H. (2012) Effect of potassium and calcium application on yield, yield components and qualitative characteristics of peanut (*Arachis hypogaea* L.) in Guilan Province, Iran. *World Appl. Sci. J.* 16(4): 540-546.

Ghasemi, S., Khoshgoftarmanesh, A.H., Afyuni, M., Hadadzadeh, H. (2014) Iron (II)-amino acid chelates alleviate salt-stress induced oxidative damages on tomato grown in nutrient solution culture. *Scientia Horticulturae*, 2014.165: 91-98

Ghasemi, S., Khoshgoftarmanesh A.H., Afyuni, M. and Hadadzadeh, H. (2013) The effectiveness of foliar applications of synthesized zinc-amino acid chelates in comparison with zinc sulfate to increase yield and grain nutritional quality of wheat. *European Journal of Agronomy*. 45: 68-74

Jones, J.B. (1999) Tomato plant culture: in the field, green house and home garden. *CRS Press, LLC Florida*. 11-53.

Kamara, E.G., Olympio, N.S and Asibuo, J.Y. (2011) Effect of calcium and phosphorus fertilizer on the growth and yield of groundnut (*Arachis hypogaea* L.). *International Research Journal of Agricultural Science and Soil Science*. 1(8).326-331.

Kamara, E.G., Olympio, N.S., JY, J.Y.A., Kabbia, M.K., Yila, K.M and Conteh, A.R. (2017) Effect of Calcium and Phosphorus Fertilizer on Seed Yield and Nutritional Quality of Groundnut (*Arachis hypogaea* L.). *International Journal of Agriculture and Forestry*. 7 (6). 129-133.

Kogram, C., Maneekao, S., Simungkhun, B and Poo Sri, B. (1999) Effect of calcium rate on the decreasing of unfilled pod of peanut (*Arachis hypogaea* L.) grown in sandy loam soil in Yasothorn province. *Field Crops Research Inst.*

Lowry, O.H., Rosebrough, A.L., Farr and Randall, R.J. (1951) *Journal of Biology and Chemistry*. 193: 265.

Madani, B., Wall, M., Mirshekari, A., Bah, A and Mohamed, M.T.M. (2015) Influence of calcium foliar fertilization on plant growth, nutrient concentrations, and fruit quality of papaya. *Hort Technology*. 25(4):496-504. Metropolitan Book Co. New Delhi, India.

Singh, A.L. (1999) Mineral Nutrition of Groundnut. In: A. Hemantranjan (ed.), *Advances in Plant Physiology*. 2:161-200. Scientific Publishers, Jodhpur, India.

Singh, A.L., Joshi, Y.C., Chaudhari Vidya and Zala, P.V. (1990) Effect of different sources of iron and sulphur on leaf chlorosis nutrient uptake and yield of ground. *Fert.Res.*24:85-96.

Son, S., Lee, E.S. and Lee, E. W. (1974) The effects of Ca and K on the growth and yield of groundnut (*Arachis hypogaea* L.). *Research Reports of the Office of Rural Development Crop.* 16: 25-31. *Field Crop Abstracts* 28: 7342, 1975.

Souri, M.K., Yarahmadi, B. (2016) Effect of amino chelates foliar application on growth and development of marigold (*Calendula officinalis*) plants. *Iranian J. of Plant Prod. Techn.*15 (2): 109-119.

Souri, M.K. (2015) Chelates and aminochelates; and their role in plant nutrition. *Agriculture Education and Extension Press*, Tehran-Iran. (In Persian).172.

Tiwari, P.N., Gambhit, P.N. and Rajan, T.S. (1974) Rapid and non destructive determination of seed oil by pulsed nuclear magnetic resonance technique. *Journal of American oil Chemical Society* .305-308.

Abbreviation: DAS- Days After Sowing; RDF - Recommended Dose of Fertilizers; Nova Ami -Ca - Nova Amino Chelated Calcium; g-grams.