

# Response of Foliar Application of Nova Amino Chelated-Boron on Vegetative Growth, Fruit Yield and Quality of Tomato

Venkata narayana, P., Sambasiva rao, Y., Kiran kumar, A., Sreekanth, Y. and Basanth, N.

Nova Innovation Centre, Nova Agri Tech Ltd., Plot No. 57, Hanuman Nagar, Chinnatokatta, New Bowenpally, Secunderabad, Telangana – 500 011

Corresponding author email id: [cro@novaagritech.com](mailto:cro@novaagritech.com)

**Abstract** – An investigation was carried out during *kharif*, 2018 in tomato crop by application of Nova Ami-B in combination with recommended dose of fertilizers to evaluate the efficiency of Nova amino chelated boron. The experiment was laid out in Randomized Block Design(RBD) with five treatments and control i.e T<sub>1</sub>: Control (No application of any fertilizer), T<sub>2</sub>: Recommended Dose of Fertilizers (RDF), T<sub>3</sub>: T<sub>2</sub>+ 2.5g/l Boron at 20 and 45 DAT, T<sub>4</sub>: T<sub>2</sub> + 2.5ml/l of Nova Ami-B at 20 and 45 DAT, T<sub>5</sub>: 2.5g/l Boron at 20 and 45 DAT, T<sub>6</sub>: 2.5ml/l of Nova Ami-B at 20 and 45 DAT. Experiment revealed that application of (T<sub>4</sub>) Recommended Dose of Fertilizers + Nova Ami-B resulted in improved plant height, number of flowers per plant, number of fruits per plant, fruit weight per plant, total soluble solids and lycopene content over other treatments but T<sub>3</sub> showed more number of leaves per plant.

**Keywords** – Amino Chelate, Boron, Plant Height, Tomato. **Abbreviations:** TSS - Total Soluble Solids; Ami-B- Amino chelated Boron; KG- Kilograms; RDF-Recommended Dose of Fertilizers; DAT- Days After Transplanting.

## I. INTRODUCTION

Tomato is one of the most important vegetable crop in the world. It ranks third in the world's vegetable production next to potato and sweet potato, placing itself first as processing crop among the vegetables. Tomato is important and popular vegetable in India. India is a second largest producer (11.5%) of tomato in the world after China. Tomato plays an important role in human nutrition by providing essential amino acids, vitamins and minerals [1]. It also contains lycopene, which is very important antioxidant and can prevent cancer.

Boron (B) is a micronutrient critical to the growth and health of all crops. Boron plays a key role in a diverse range of plant functions including cell wall formation, stability, maintenance of structural and functional integrity of biological membranes. Reported that Boron is involved in movement of sugar or energy into growing parts of plants, pollination and seed set [2]. Have suggested the involvement of boron in stomatal regulation [3].

In crop production boron (B) is one of the essential micronutrient required for normal growth of most of the crops. Boron fertilization is necessary for improvement of crop yield as well as nutritional quality. Boron fertilization improves photosynthetic activity, enhances activity of enzymes and plays significant role in protein and nucleic acid metabolism [4]. The boron requirement is much higher for reproductive growth than for vegetative growth in most plant species. Boron increases flower production and

retention, pollen tube elongation and germination, seed and fruit development. Foliar application of boron is preferred over soil application because of the very narrow range from deficient to toxic levels [5].

## II. MATERIAL AND METHODS

A field experiment on tomato (cv. Pusa Ruby) was conducted in the *kharif*, 2018 at Nova Agritech Ltd. Experimental farm, Kamareddy, Telangana, India. To determine the effect of Nova amino chelated-boron on yield components and quality parameters of tomato. The six treatments were replicated in four times in a randomized block design. The field was prepared by ploughing and leveling, divided into small plots of 25m<sup>2</sup> (5m×5m) and the tomato seedlings were raised on nursery beds. Then 25 days after, seedling with 3-4 leaves transplanted in prepared plots with a distance of 60cm×60cm within and between the rows. The crop was grown under recommended package of practices and proper care was taken to protect it from weeds, insects, pests and diseases during entire cropping season. Data was recorded on established plants at 10 days after sprayings. Plant height, number of leaves per plant, number of flowers per plant, number of fruits per plant, fruit weight, lycopene content, total soluble sugars and yields were recorded. A total of two foliar sprays were carried out during the cropping season.

Table 1. Treatments:

T <sub>1</sub>	Control (No application of any fertilizer).
T <sub>2</sub>	Recommended dose of fertilizers (RDF)
T <sub>3</sub>	T <sub>2</sub> + 2.5g/l Boron at 20 and 45 DAT
T <sub>4</sub>	T <sub>2</sub> + 2.5ml/l of Nova Ami-B at 20and 45 DAT.
T <sub>5</sub>	2.5g/l Boron at 20 and 45 DAT
T <sub>6</sub>	2.5ml/l of Nova Ami-B at 20and 45 DAT

## III. RESULT AND DISCUSSIONS

The findings in the present study as well as relevant discussion have been presented under following heads.

Table 2. Effect of Nova Ami-Boron on Growth and Yield of Tomato.

Treatments	Plant height (cm)	No. of leaves/plant	No. of flowers /plant	No. of fruits /plant	Fruit weight/ Plant (kg)	TSS (°Brix)	Lycopene content mg/100g
T <sub>1</sub>	85.00	55.50	35.75	18.75	0.710	3.69	4.408
T <sub>2</sub>	110.50	67.00	76.75	53.25	1.248	5.38	4.798
T <sub>3</sub>	116.75	75.50	90.25	56.25	1.748	5.53	4.845
T <sub>4</sub>	121.88	74.75	95.75	67.25	1.903	5.64	4.918
T <sub>5</sub>	99.75	58.50	66.50	34.50	1.003	4.54	4.600
T <sub>6</sub>	107.5	60.00	61.50	40.50	1.208	4.70	4.683
SEm	2.1755	2.5275	3.3019	1.8981	0.0510	0.0790	0.0816
CD 5%	6.5575	7.618	9.9531	5.7215	0.1538	0.2382	0.2458
CV	4.07	7.752	9.2903	8.4204	7.8345	3.2155	3.4641

**A. Plant Height**

Plant height ranged from 122 to 85 cm. Among the various treatments, application of RDF + 2.5ml/l of Nova Ami-B at 20 and 45 DAT (T<sub>4</sub>) results significantly higher plant height of 122cm. Reported that higher dose of boron and medium dose of nitrogen were influential nutrients for increasing the plant height [6]. Boron increased plant height and the number of branches of tomato plant by promoting root growth, which enhances nutrient absorption [7]. Deficiency of calcium and boron decreases plant height by decreasing mitotic activity in the terminal meristem [8].

**B. Number of Leaves per Plant**

Results revealed that significant differences were present in number of leaves per plant. Among six treatments T<sub>3</sub> (T<sub>2</sub> + 2.5g/l Boron at 20 and 45 DAT) recorded the highest number of leaves followed by T<sub>4</sub> (T<sub>2</sub> + 2.5ml/l of Nova Ami-B at 20 and 45 DAT). Boron enhance the number of metabolites necessary for building plant organs, consequently the vegetative growth of plants [9]. The foliar application of B and Zn have a significant effect on overall development of plant [10].




Fig. 1. Influence of Nova Ami-B on plant height, no. of leaves, no. of flowers and no. of fruits.




Fig. 2. Influence of Nova Ami-B on Fruit weight, TSS and Lycopene content.

### C. Number of Flowers per Plant

The present research work showed that maximum number of flowers observed in T<sub>4</sub> (T<sub>2</sub> + 2.5ml/l of Nova Ami-B at 20 and 45 DAT) and minimum number of flowers recorded in T<sub>1</sub> (Control). Stated that higher dose of boron favored the higher number of flower clusters in tomato [6]. Reported that optimum supply of boron promoted more flower clusters formation [12]. Foliar application of borax resulted in a significant increase in the number of flowers per cluster [11].

### D. Number of Fruits per Plant

This observation indicates that Highest number of fruits found in T<sub>4</sub> which differed significantly with all other treatments. Lowest number of fruits recorded in T<sub>1</sub> (Control). Observed that number of fruits per cluster was increased with increasing level of boron [6]. The application of boron enhances fruit set [13] by delaying abscission of flowers [14]. These results agree with reported by [15].

### E. Fruit Weight per Plant

The obtained results shows fruit weight per plant varied from 1.903 to 0.710. Maximum fruit weight was observed in T<sub>4</sub> which shows significant differences with other treatments. Minimum fruit weight was recorded in T<sub>1</sub>. The fruit weight of tomato increased significantly with foliar application of boron [11].

### F. Total Soluble Solids:

The present study confirms that maximum TSS was recorded in T<sub>4</sub> which differed significantly with T<sub>1</sub>, T<sub>2</sub>, T<sub>5</sub>, T<sub>6</sub> and showed parity with T<sub>3</sub>. The total soluble solids (TSS) in tomato fruit was extremely low at the lowest B level. B is needed for the production of uracil, a pre-cursor of uridine-diphosphate-glucose (UDPG), an essential enzyme for the production of sucrose. Sucrose is easily translocated, and important for carbohydrate translocation [14].

### G. Lycopene Content

The present study revealed that significant differences were observed among treatments. Maximum amount of lycopene content observed in T<sub>4</sub> which shows on par with T<sub>3</sub>, T<sub>2</sub> and minimum amount of lycopene recorded in T<sub>1</sub>. Lycopene content of the fruit gradually increased with the increasing boron [16].

## IV. CONCLUSION

It can be concluded that from the study application of Recommended Dose of Fertilizers + Nova Ami-B resulted in improved plant height, number of flowers per plant, number of fruits per plant, fruit weight per plant, total soluble solids and lycopene content over other treatments.

## REFERENCES

- [1] Sainju, U., Dris, R and Singh, B. 2003. Mineral nutrition of tomato. *Journal of Food Agriculture and Environment*. 2 (1): 176-183.
- [2] Parr, A.J., Loughmann, B.C. 1983 Metals and Micronutrients Uptake and Utilization by Plants. *Academic Press*, New York. USA. 87-107.
- [3] Sharma P.N and Tanuja, R. 1991 *Indian Journal of Experimental Biology*. 29(10): 967-970.

- [4] Kolesnik, L. V. 1962. *Proc. Kishinev. Agri. Inst.* 27: 3-9.
- [5] Reisenauer, H.M., Walsh, L.M and Hoefl, R.G. 1973 testing soils for sulfur, boron, molybdenum, and chlorine, In: *Soil Testing and Plant Analysis* (Eds. LM Walsh, JD Benton). *Soil Science Society of America*. Madison, Wisconsin. 173.
- [6] Haque, M.E., Paul, A.K and Sarker, J.R. 2011. Effect of nitrogen and boron on the growth and yield of tomato (*Lycopersicon esculentum* M.). *Intern. J. Bioresource Stress Manag.* 2(3):277-282.
- [7] Sathya, S., Mani, S., Mahedran, P.P and Arulmozhiselvan, K. 2010. Effect of application of boron on growth, quality and fruit yield of PKM 1 tomato. *Indian J Ag Res.* 44: 274-280.
- [8] Nelson, P.V., Niedziela, C.E. 1998 Effect of calcium sources and temperature regimes on calcium deficiency during hydroponic forcing of tulip. *Sci Hortic.* 73: 137-150.
- [9] Marschner, H. 1995. The soil root interface (rhizosphere) in relation to mineral nutrition. *Mineral nutrition of higher plants*.
- [10] Sultana, S., Naser, H.M., Akhter, S and Begum, R.A. 2016. Effectiveness of soil and foliar applications of zinc and boron on the yield of tomato. *Bang. J of Agr Res* 41(3): 411-418.
- [11] Rab, A and Haq, I.U. 2012. Foliar application of calcium chloride and borax influences plant growth, yield, and quality of tomato (*Lycopersicon esculentum* Mill.) fruit. *Turkish Journal of Agriculture and Forestry*. 36(6): 695-701.
- [12] Day, S.C. 2000. Tomato crop in vegetable growing. *Agrobios*, New Delhi, India. 59-61.
- [13] Desouky, I.M., Haggag, L.F., El-Migeed, M.A., Kishk, Y.F.M. and El-Hady, E.S., 2009. Effect of boron and calcium nutrients sprays on fruit set, oil content and oil quality of some olive oil cultivars. *World Journal of Agricultural Sciences*. 5(2):180-185.
- [14] Smit, J.N and Combrink, N.J.J. 2004. The effect of boron levels in nutrient solutions on fruit production and quality of greenhouse tomatoes. *South African Journal of Plant and Soil*. 21(3): 188-191.
- [15] El-Shall M.A, El-Araby S.M, Ghoneim I.M, Anter, H. (2003) Effect of bio fertilization under varying NPK levels on growth, yield and fruit quality of strawberry plants. *J Agri Env Sci.* Alex Univ. 2: 106-129.
- [16] Salam, M.A., Siddique, M.A., Rahim, M.A., Rahman, M.A and Goffar, M.A. 2011. Quality of tomato as influenced by boron and zinc in presence of different doses of cowdung. *Bangladesh Journal of Agricultural Research*. 36(1).151-163.

## AUTHOR'S PROFILE



Mr. Venkata narayan P. M.Sc (Agriculture) with NET studied in ANGRA University and present working as a research associate at the department of Research and development of NOVA AGRITECH LIMITED. He has published 6 research articles and 10 popular articles. Mr. Venkata narayana work has been published in some journals like Int. J. Curr. Microbiol. App. Sci, International Journal of Agricultural Science and Research (IJASR), international journal of science and nature etc. He is a member of plant physiology club.